ILLINOIS AT URBANA-CHAMPAIGN Institute of Natural Resource Sustainability William W. Shilts, Executive Director ILLINOIS STATE GEOLOGICAL SURVEY E. Donald McKay III, Interim Director For more information contact: Institute of Natural Resource Sustainablity Illinois State Geological Survey 615 East Peabody Drive Champaign, Illinois 61820-6964

(217) 333-4747

http://www.isgs.illinois.edu

## Danville Coal Sulfur EFFINGHAM County


County Coal Map Series Andrew Louchios, Scott Elrick, Chris Korose, David Morse


Map construction: October 29, 2009

This product is under review and may not meet the standards of the Illinois State Geological Survey.

County coal maps and select quadrangle maps available as downloadable PDF files at: http://www.isgs.illinois.edu/maps-data-pub/coal-maps/county-index.shtml







western Kentucky. The coal crops out along the margins of the Illinois Basin and reaches a maximum depth in Illinois of about 1,200 feet. (See Fig 1. and Fig 2.) The Danville Coal is in the Shelburn formation which is part of the Desmoninesian Series. The Danville Coal has been mined in Livingston, McLean, La Salle, and Marshall Counties in addition to Vermilion County. In most of the remainder of the state it is a thin coal, generally a few inches to less than 3 feet thick. The Danville

Coal is generally overlain by the Farmington Shale Member of the Shelburn Formation, but in places the immediate roof is 1-2 feet of black fissile shale. It is underlain by a kins, 1968 - B95). (See Fig 4.)

The original resource of Dan-

Avail. w/ potential restr. 12.5 208.6 mined. Approximately 23% of the original Danville Coal resources, 4.5 billion the type locality in Vermillion tons, are considered available for mining. (See Fig 3.) Available means that to be too thin or too county, the Danville Coal is 6 the surface land-use and geologic conditions related to mining of the deposit poor in quality to jusfeet thick and occurs 20 feet (e.g. thickness, depth, in-place tonnage, stability of bedrock overburden) are above the Herrin Coal. (Hop-comparable to other coals currently being mined in the state. Of these resources, 4 billion tons occur in coal 42 to 66 inches thick and 0.4 billion tons occur in thicknesses greater than 66 inches.

mining was in east-central Illinois near the city of Danville where the coal has

ville Coal in the State of Illi- **T**he Danville Coal has been mined in Illinois for over 100 years, but only about

ground methods. Except for mines in east-central Illinois, most large surface mines recover the Danville Coal only as part of their operation to remove overburden to mine the underlying Herrin Coal. In many cases, the Danville seam

Fig. 4 Pennsylvanian Stratigraphic Column					
Series	Fm.	Graphic Column	Central and Southern Members and Beds	Northern and Western Members and Beds	Eastern and Southern Members and Beds
			Trivoli Sandstone  Scottville Limestone Athensville Coal (SW)	Trivoli Sandstone Exline Limestone	Trivoli Sandstone
Desmoinesian Series	ale Shelburn	****	Lake Creek Coal Pond Creek Coal Gimlet Sandstone Rock Branch (SW)/ DeGraff (S) Coal	Lonsdale Limestone Gimlet Sandstone	West Franklin Limestone
		*******	Piasa Limestone  Danville Coal  Galum Limestone  Allenby Coal  Bankston Fork Limestone	Farmington Shale Danville Coal	Danville Coal  Bankston Fork Limestone
			Anvil Rock Sandstone Conant Limestone Jamestown Coal	Copperas Creek Sandstone Lawson Shale	Anvil Rock Sandstone  — Conant Limestone  — Jamestown Coal
			Brereton Limestone Anna Shale Energy Shale	Brereton Limestone Anna Shale	Brereton Limestone Anna Shale
	Carbondale	***********	Herrin Coal	Herrin Coal Spring Lake Coal Bed Big Creek Sandstone Vermillionville Sandstone	Herrin Coal
	Ü	XXXXXXX	Briar Hill Coal		Briar Hill Coal

**Coal Sulfur** 

Less than or equal to 0.40 (lb S per MM Btu)

0.41 to 0.60 (lb S per MM Btu)

0.61 to 0.83 (lb S per MM Btu)

0.84 to 1.24 (lb S per MM Btu)

1.25 to 1.67 (lb S per MM Btu)

1.68 to 2.50 (lb S per MM Btu)

Greater than 2.50 (lb S per MM Btu)

tify recovery and was simply discarded in the spoil pile with other rock overburden. (Modified from ISGS Pub. IM 124, Korose, et al)

- Handbook of Illinois Stratigraphy, 1975, Illinois State Geological Survey Bulletin 95, 261p. - Christopher P. Korose, Colin G. Treworgy, Russell J. Jacobson, and Scott D. Elrick, 2002, Availabilnois totals 19.6 billion tons, 1% of the original resource has been depleted. The most extensive area of ity of the Danville, Jamestown, Dekoven, Davis, and Seelyville Coals for mining in Selected Areas of Illinois: Illinois State Geological Survey Illinois Minerals 124, 44 p.

The maps and digital files of this study were compiled from data from a variety of public and private sources and have varying degrees of completeness and accuracy. They present interpretations of the geology of the area and are based on available data. However, these interpretations are based on data that may vary with respect to accuracy of geographic location, type, quantity, and reliability, as they were supplied to the Illinois State Geological Survey. Consequently, the accuracy of the interpreted features shown in these files is subject to the limitations of the data and varies from place to place.

Contoured features less than 7 million square feet (about 1/2 mile square) in area may not be accurately portrayed or resolved. This data set provides a large-scale conceptual model of the geology of the area on which to base further work. These data are not intended for use in site-specific screening or decision-making.

## Disclaimer

The Illinois State Geological Survey and the University of Illinois make no guarantee, expressed or implied, regarding the correctness of the interpretations presented in this data set and accept no liability for the consequences of decisions made by others on the

basis of the information presented here. © 2009 Board of Trustees of the University of Illinois. All rights reserved.