(217) 333-4747

http://www.isgs.illinois.edu

## Herrin Coal Chlorine CALHOUN County


County Coal Map Series Andrew Louchios, Scott Elrick,


Chris Korose, David Morse Map construction: October 29, 2009

This product is under review and may not meet the standards of the Illinois State Geological Survey.

County coal maps and select quadrangle maps available as downloadable PDF files at: http://www.isgs.illinois.edu/maps-data-pub/coal-maps/county-index.shtml

12N 12W 75 6W 12N 11W PIKE 7S 5W 75 4W 7S 2W 11N\_13W 11N 12W 11N 11W 14W 8\$ 2W 10N\_11W 10N 12W Ca 2010N 13W 19S 3W11 GREENE 7\_9\$ 2W 9N 1<mark>2W</mark> 9N 13W 10S 3W 10S-2W \14W L 8N 13W ∕8N 12W 3 \ \8N \11\W 11S 2W **JERSEY** CALHOUN 7N-12W 12S 12S 2W 6N 12W 12**Ş** 1V 13\$ 2W4 Eagle






orth-south cross section of the Pennsylvanian System in Illinois

The Herrin Coal underlies about two thirds of Illinois as well as portions of western Indiana and western Kentucky. The coal crops out along the margins of the Illinois Basin and reaches a maximum depth in Illinois of about 1,300 feet. (See Fig . and Fig 2.) The Herrin Coal is a normal brightbanded coal. Its lower portion contains a prominent claystone parting (the "blue band") that normally is 1-3 inches thick. It averages more than 6 feet thick in extensive areas and locally reaches 15 feet. It is thin in much of central Illinois but has

been extensively mined in western, west-central, southern, and northern Illinois, as well as in the southern part of the Danville region of eastern Illinois. In some places the coal is cut out by channels filled with the Anvil Rock much as a mile wide and 60-



Sandstone Member. In parts is overlain by relatively thick bodies of the gray shale of up to a few tens of greater than 66 inchof Illinois, silty gray shale as feet it has a much lower sulfur content than elsewhere. The gray shale over- es. (Modified from much as 100 feet thick over- lies the coal principally in parts of Williamson, Franklin, Jefferson, Madison, ISGS Pub. IM 120, lies the Herrin Coal. Associ- St. Clair, eastern Macoupin, and S. Vermilion. Generally, however the Herrin Treworgy, et al) ated with this shale is a chan- Coal is overlain by either the Anna Shale Member (black fissile shale) or the nel sandstone commonly as Brereton Limestone Member. (Hopkins, 1968 - B95, See Fig 4.)

80 feet thick mapped as Anvil The original resource of Herrin Coal in the State of Illinois totals 88.5 billion References: Rock Sandstone and may be tons, of which 9.4 billion have been mined. Approximately 58% of the original - Handbook of Illinois Stratigraphy, 1975, Illinois State Geological Survey Bulletin 95, 261p. contemporaneous with the Herrin Coal resources, 51 billion tons, are considered available for mining. (See - Treworgy, C.G., C.P. Korose, C.A. Chenoweth, and D.L. North, 1999a, Availability of the Herrin coal. In areas where the coal Fig 3.) Available means that the surface land-use and geologic conditions re-Coal for mining in Illinois: Illinois State Geological Survey Illinois Minerals 120, 54 p.

Pennsylvanian Stratigraphic Column

Central and Southern Northern and Western Eastern and Southern Members and Beds Members and Beds Members and Beds

peras Creek Sandstone Lawson Shale

Brereton Limestone Anna Shale

St. David Limestone Turner Mine Shale

Hanover Limestone Excello Shale Houchin Creek Coal Breezy Hill Limestone Kerfon Creek Coal Pleasantview Sandstore

Anvil Rock Sandston

rereton Limesto Anna Shale

lated to mining of the Fig. 4

deposit (e.g. thick-

ness, depth, in-place

tonnage, stability

of bedrock overbur-

den) are comparable

to other coals cur-

rently being mined

in the state. Of these

resources, 21 billion

42 to 66 inches thick

and 30 billion tons

occur in thicknesses

## **Coal Chlorine** Less than 0.1 % 0.1 to 0.2 % 0.2 to 0.3 % 0.3 to 0.4 % 0.4 to 0.5 % 0.5 to 0.6 % Greater than 0.6 %

## **Map Explanation**

The maps and digital files of this study were compiled from data from a variety of public and private sources and have varying degrees of completeness and accuracy. They present interpretations of the geology of the area and are based on available data. However, these interpretations are based on data that may vary with respect to accuracy of geographic location, type, quantity, and reliability, as they were supplied to the Illinois State Geological Survey. Consequently, the accuracy of the interpreted features shown in these files is subject to the limitations of the data and varies from place to place.

Contoured features less than 7 million square feet (about 1/2 mile square) in area may not be accurately portrayed or resolved. This data set provides a large-scale conceptual model of the geology of the area on which to base further work. These data are not intended for use in site-specific screening or decision-making.

## **Disclaimer**

The Illinois State Geological Survey and the University of Illinois make no guarantee, expressed or implied, regarding the correctness of the interpretations presented in this data set and accept no liability for the consequences of decisions made by others on the

basis of the information presented here.  $\hbox{@}$  2009 Board of Trustees of the University of Illinois. All rights reserved.