LLINOIS Institute of Natural Resource Sustainability William W. Shilts, Executive Director **ILLINOIS STATE GEOLOGICAL SURVEY** E. Donald McKay III, Interim Director For more information contact: Institute of Natural Resource Sustainablity

Illinois State Geological Survey

Champaign, Illinois 61820-6964

615 East Peabody Drive

http://www.isgs.illinois.edu

(217) 333-4747


## Herrin Coal Chlorine WAYNE County


County Coal Map Series Andrew Louchios, Scott Elrick,


> Chris Korose, David Morse Map construction: October 29, 2009

This product is under review and may not meet the standards of the Illinois State Geological Survey.

County coal maps and select quadrangle maps available as downloadable PDF files at: http://www.isgs.illinois.edu/maps-data-pub/coal-maps/county-index.shtml







orth-south cross section of the Pennsylvanian System in Illinois

The Herrin Coal underlies about two thirds of Illinois as well as portions of western Indiana and western Kentucky. The coal crops out along the margins of the Illinois Basin and reaches a maximum depth in Illinois of about 1,300 feet. (See Fig . and Fig 2.) The Herrin Coal is a normal brightbanded coal. Its lower portion contains a prominent claystone parting (the "blue band") that normally is 1-3 inches thick. It averages more than 6 feet thick in extensive areas and locally reaches 15 feet. It is thin in much of central Illinois but has been extensively mined in western, west-central,

southern, and northern Illinois, as well as in the southern part of the Danville region of eastern Illinois. In some places the coal is cut out by channels filled with the Anvil Rock Sandstone Member. In parts is much as a mile wide and 60-

				Original	Mined	R emaining	Available
Danville				19.6	0.2	19.4	4.5
Jamestown				3.6	0	3.6	0.9
Herrin				88.5	9.4	79.0	51.0
Springfield				65.1	2.2	63.0	27.0
Colchester				19.0	0.5	18.5	1.0
Dekoven	]			6.0	0.1	5.9	0.3
Davis		■ Available ■ Avail. w/ poten	itial restr.	9.6	0.1	9.5	4.7
Seelyville		■ Restricted or n	nined	9.7	0	9.7	6.7
0	20	40 60	80 1	100	(All number	s in Billions of	f Tons)
Fig. 3		billions of tons		221.1	12.5	208.6	96.1
s overlain by relatively thick bodies of the gray shale of up to a few tens of							

of Illinois, silty gray shale as feet it has a much lower sulfur content than elsewhere. The gray shale over- es. (Modified from much as 100 feet thick over- lies the coal principally in parts of Williamson, Franklin, Jefferson, Madison, lies the Herrin Coal. Associ- St. Clair, eastern Macoupin, and S. Vermilion. Generally, however the Herrin Treworgy, et al) ated with this shale is a chan- Coal is overlain by either the Anna Shale Member (black fissile shale) or the nel sandstone commonly as Brereton Limestone Member. (Hopkins, 1968 - B95, See Fig 4.)

occur in thicknesses greater than 66 inch-ISGS Pub. IM 120,

deposit (e.g. thick-

ness, depth, in-place tonnage, stability

of bedrock overbur-

den) are comparable

to other coals cur-

rently being mined

in the state. Of these

resources, 21 billion

tons occur in coal

42 to 66 inches thick

and 30 billion tons

80 feet thick mapped as Anvil The original resource of Herrin Coal in the State of Illinois totals 88.5 billion References: Rock Sandstone and may be tons, of which 9.4 billion have been mined. Approximately 58% of the original - Handbook of Illinois Stratigraphy, 1975, Illinois State Geological Survey Bulletin 95, 261p. contemporaneous with the Herrin Coal resources, 51 billion tons, are considered available for mining. (See - Treworgy, C.G., C.P. Korose, C.A. Chenoweth, and D.L. North, 1999a, Availability of the Herrin coal. In areas where the coal Fig 3.) Available means that the surface land-use and geologic conditions re-Coal for mining in Illinois: Illinois State Geological Survey Illinois Minerals 120, 54 p.


Central and Southern Northern and Western Eastern and Southern Members and Beds Members and Beds Members and Beds

St. David Limestone Turner Mine Shale

Hanover Limeston Excello Shale

Anvil Rock Sandston

Brereton Limesto Anna Shale



## **Map Explanation**

The maps and digital files of this study were compiled from data from a variety of public and private sources and have varying degrees of completeness and accuracy. They present interpretations of the geology of the area and are based on available data. However, these interpretations are based on data that may vary with respect to accuracy of geographic location, type, quantity, and reliability, as they were supplied to the Illinois State Geological Survey. Consequently, the accuracy of the interpreted features shown in these files is subject to the limitations of the data and varies from place to place.

Contoured features less than 7 million square feet (about 1/2 mile square) in area may not be accurately portrayed or resolved. This data set provides a large-scale conceptual model of the geology of the area on which to base further work. These data are not intended for use in site-specific screening or decision-making.

## Disclaimer

The Illinois State Geological Survey and the University of Illinois make no guarantee, expressed or implied, regarding the correctness of the interpretations presented in this data set and accept no liability for the consequences of decisions made by others on the basis of the information presented here.

© 2009 Board of Trustees of the University of Illinois. All rights reserved.